Non-Irrigated Recharge Experiment

Allan Wylie
IDWR

Outline

- Changes to non-irrigated recharge
- Physical properties
- Steady state match
  - Head
  - Gain and loss
- Transient match
  - Head
  - Gain and loss match

Changes to non-irrigated recharge



- None (basalt) - 1.74
- Thin soil – 0.14
- Thick soil – 1e-10
- Other – not adjusted

Transmissivity





Storage


Legend
Sy
Unitless
High : 0.279606
Low : 0.005172
seas.txt Events
masmeas.txt Events


Legend
SY pilot points
1X1_cell_attribs
sy
0.000 - 0.010
0.011 - 0.050
0.051 - 0.100
0.101 - 0.150
0.151 - 0.380

Riverbed conductance





Drain Conductance





Steady state



- Head
  - Pink line is 45°
  - If this were a perfect match
    - Y=1x + 0
    - $R^2 = 1$

Steady state



Steady state



- Gain and loss match
  - Pink line is 45°
  - If this were a perfect match
    - Y=1x + 0
    - $R^2 = 1$

Seasonal data set



- Hundreds to thousands of observations
  - Lots of data
  - Limited spatial coverage

Seasona


Legend
AbsMean (ft)
0.546930 - 1.043700
1.043701 - 1.453400
1.453401 - 2.309600
2.309601 - 3.078400
3.078401 - 5.338700


771
Feet
18
16
14
12
10
8
6
4
2
0
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

Feet
18
16
14
12
10
8
6
4
2
0
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

Seasonal


Legend
AbsMean (ft)
0.546930 - 1.04370
1.043701 - 1.45340
1.453401 - 2.30960
2.309601 - 3.07840
3.078401 - 5.33870

Feet
16
14
12
10
8
6
4
2
0
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


282
Feet
14
12
10
8
6
4
2
0
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

data set


942
20
10
0
-10
-20
-30
-40
-50
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


3.078401 - 5.338700

942
0
-10
-20
-30
-40
-50
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

Seasonal data set



Mass measurement



- Observations collected on behalf of this modeling effort
  - 3/1/2001 –5/8/2002

Mass measurement







Mass measurement



Mass measurement



Mass measurement



Trend



- Annual observations in wells

d


327
Feet
10
9
8
7
6
5
4
3
2
1
0
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


327
Feet
6
5
4
3
2
1
0
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


1062
Feet
5
4
3
2
1
0
-1
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled
Feet
4
3
2
1
0
-1
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

d


1253
Feet
20
15
10
5
0
-5
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


1253
Feet
5
0
-5
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

Trend


3386
Feet
14
12
10
8
6
4
2
0
-2
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


3386
Feet
16
14
12
10
8
6
4
2
0
-2
-4
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measu
Model



Trend


Legend
Trend AbsMean (ft)
0.023000 - 1.891000
1.891001 - 4.421600
4.421601 - 8.498500
8.498501 - 14.049000
14.049001 - 27.758000


4213
Feet
7
6
5
4
3
2
1
0
-1
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


4213
Feet
7
6
5
4
3
2
1
0
-1
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled



Snake River



- Snake River reaches


Ashton-Rexberg
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
100.0
80.0
60.0
40.0
20.0
0.0
-20.0
-40.0
May-80
Jan-83
Oct-85
Jul-88
Apr-91
Jan-94
Oct-96
Jul-99
Mar-02
Monthly reach gain or loss
3NonIrrRch
Monthly reach gain or loss
Modeled


Hiese-Shelly
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
100.0
50.0
0.0
-50.0
-100.0
-150.0
May-80
Jan-83
Oct-85
Jul-88
Apr-91
Jan-94
Oct-96
Jul-99
Mar-02
Monthly reach gain or loss
3NonIrrRch
Total Monthly Reach Gain (+) or Loss (-, In Thousands of AC-FT Per Month
Monthly reach gain or loss
Modeled


Shelly-nr Blackfoot
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
150.0
100.0
50.0
0.0
-50.0
-100.0
-150.0
-200.0
May-80
Jan-83
Oct-85
Jul-88
Apr-91
Jan-94
Oct-96
Jul-99
Mar-02
Monthly reach gain or loss
3NonIrrRch
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
Monthly reach gain or loss
Modeled


nr Blackfoot-Neeley
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
300.0
250.0
200.0
150.0
100.0
50.0
0.0
-50.0
May-80
Jan-83
Oct-85
Jul-88
Apr-91
Jan-94
Oct-96
Jul-99
Mar-02
Monthly reach gain or loss
3NonIrrRch
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
Monthly reach gain or loss
Modeled


Neeley-Minidoka
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
150.0
100.0
50.0
0.0
-50.0
-100.0
May-80
Jan-83
Oct-85
Jul-88
Apr-91
Jan-94
Oct-96
Jul-99
Mar-02
Monthly reach gain or loss
3NonIrrRch
Total Monthly Reach Gain (+) or Loss (-), In Thousands of AC-FT Per Month
Monthly reach gain or loss
Modeled

Springs


Eastern Snake Plain
Aquifer Model: Run
Modeled River
modeledSpgs

s Washbowl


DEVILSWB
Feet
0
-200000
-400000
-600000
-800000
-1000000
-1200000
-1400000
-1600000
-1800000
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled


Feet
-600000
-800000
-1000000
-1200000
-1400000
-1600000
-1800000
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled


DEVILSC
0
-500000
-1000000
-1500000
-2000000
-2500000
-3000000
-3500000
-4000000
-4500000
-5000000
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled

vils Corral


-2500000
-3000000
-3500000
-4000000
-4500000
-5000000
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled

e Lakes


BLUELK
0.00E+00
-5.00E+06
-1.00E+07
-1.50E+07
-2.00E+07
-2.50E+07
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


-1.00E+07
-1.10E+07
-1.20E+07
-1.30E+07
-1.40E+07
-1.50E+07
-1.60E+07
-1.70E+07
-1.80E+07
-1.90E+07
-2.00E+07
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

stal Springs





r Lakes


CLEARLK
0.00E+00
-5.00E+06
-1.00E+07
-1.50E+07
-2.00E+07
-2.50E+07
-3.00E+07
-3.50E+07
-4.00E+07
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled


-2.50E+07
-2.70E+07
-2.90E+07
-3.10E+07
-3.30E+07
-3.50E+07
-3.70E+07
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

gs Spring


BRIGGS
0
-2000000
-4000000
-6000000
-8000000
-10000000
-12000000
-14000000
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled


-8000000
-8500000
-9000000
-9500000
-10000000
-10500000
-11000000
-11500000
-12000000
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Measured
Modelled



ox Canyon Spring



sand Springs






MALAD
0.00E+00
-2.00E+07
-4.00E+07
-6.00E+07
-8.00E+07
-1.00E+08
-1.20E+08
-1.40E+08
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

ad


-8.00E+07
-8.50E+07
-9.00E+07
-9.50E+07
-1.00E+08
-1.05E+08
-1.10E+08
-1.15E+08
-1.20E+08
Feet
May-80
Oct-85
Apr-91
Oct-96
Mar-02
Sep-07
Measured
Modelled

End