**Appendix A**

Description of geologic map units from Thousand Springs Quadrangle map by Gillerman et al, 2005 Idaho Geological Survey.

# DESCRIPTION OF MAP UNITS

## ARTIFICIAL DEPOSITS

m — **Made ground (Holocene)**—Artificial fills composed of excavated, transported, and emplaced construction materials typically derived locally. Primarily areas modified for fish ponds.

## ALLUVIAL AND LACUSTRINE DEPOSITS

Qam — **Alluvium of mainstreams (Holocene)**—Channel and flood-plain deposits of the Snake and Malad rivers. Stratified silt, sand, and gravel of channel bars, islands, and shorelines. Typically 1-10 feet thick.

Qoam — **Older alluvium of mainstreams (Holocene)**—Channel and flood-plain deposits of the Snake River that form fill terraces 10-50 feet above reservoir level. Primarily beds of sand and pebble gravel overlain by bedded to massive silt and sand. Grades and interfingers laterally into colluvium and alluvial-fan deposits along valley sides. The Buckeye Ranch terrace west of Hagerman is anomalously high at 2,850 feet elevation, about 50 feet above the reservoir. The aggradation may represent response to a higher base-level caused by landslides downstream.

Qaf — **Alluvial-fan deposits (Holocene)**—Stratified silt, sand, and minor gravel that form alluvial fans at the base of steep canyon slopes. Fans modify landslide deposits in the Snake River canyon west of Hagerman. North of Lower Salmon Falls Dam alluvial-fan deposits merge and interstratify with older mainstream alluvium (*Qoam*). Thickness probably 5-30 feet.

Qas — **Alluvium of side streams (Holocene)**—Channel and flood-plain deposits of Yahoo Creek, Billingsley Creek, and minor tributaries to the Snake River. Primarily stratified silt, sand, and minor gravel.

Qoab

**Older alluvium of Big Wood River (Holocene or Pleistocene)**—Cobble and boulder gravel deposited by high-energy floods of the Big Wood River (Malad River). Gravel forms four terraces (*Qoab*[1] - *Qoab*[4]) that are 10 feet, 25 feet, 50 feet, and 100 feet, respectively, above the present river level.

Qams

**Scoured flood pathways of side streams (Holocene or Pleistocene)**—Surface of basalt scoured by one or more Malad-River overland floods from the Big Wood River that augmented headward erosion of Malad Gorge. Common small scour marks on basalt surfaces are aligned with the direction of water flow. Surface is mantled with thin and discontinuous sand and gravel deposits.

**Bonneville Flood Deposits**

Qabg

**Sand and gravel in giant flood bars (Pleistocene)**—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlined giant expansion bars with large-scale crossbeds. Deposited during highest-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qabgl

**Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)**—Stratified coarse sand to sandy pebble-cobble gravel deposited in eddies, side-channel positions, and lower-energy, waning-stage flood channels.

Qabs

**Scabland of flood pathways (Pleistocene)**—Flood-scoured surface with variable amounts of flood deposits. In the Hagerman Valley sedimentary cover has been stripped and basalt surfaces have been plucked, gouged, and smoothed. On sedimentary units such as Yahoo Clay, flood waters downcut, backcut, channeled, and steamlined the pre-flood surface. Includes minor deposits of coarse sand that are not mapped at this scale.

Qy

**Yahoo Clay (Pleistocene)**—Laminated to thin-bedded clay and silty clay. Pinkish white to light yellowish brown and conchoidal fracture when dry. Common partings along bedding and vertical jointing produce small blocks when exposed. Malde (1982) described the type locality near the mouth of Yahoo Creek, the lava-dam origin, and the distribution of the clay in the Snake River canyon from near Bliss to the Melon Valley. Stratigraphic evidence demonstrates the Yahoo Clay is younger than the basalt of Notch Butte (*Qnb*), but older than the Bonneville Flood. Malde (1982) attributes the clay to McKinney Lake, a temporary lake formed by damming of the Snake River by basalt of McKinney Butte (*Qmk*). Malde's interpretation of the lake is compelling and his stratigraphic evidence was confirmed by our field mapping. East and south of Hagerman, Malde and Powers (1972) and Covington and Weaver (1990) show the Yahoo Clay buried by Crowsnest Gravel except where dissected. However, our field mapping and the soil survey by Johnson (2002) suggest the Yahoo Clay is the significant mappable unit at the land surface. Yahoo Clay was scoured by the maximum stage of the Bonneville Flood. Flood features include streamlined topography and a relict cataract and plunge pool one-half mile east of Hagerman.

Tt

**Tuana Gravel (Pliocene)**—Well-bedded and sorted pebble and cobble gravel interbedded with layers of sand, silt, and clay. Gravel clasts are well rounded and commonly disc-shaped. Imbrication of clasts is common. Gravel-clast lithologies and imbrication directions suggest the gravel was deposited by an ancestral Salmon Falls Creek that prograded braided-stream deposits across a high, nearly flat plain formed on the Glenns Ferry Formation. Original extent unknown owing to erosion. Highest erosion remnants are mantled by 10-25 feet of loess with several buried soils. A thick duripan just above the gravel often forms an erosion-resistant cap "rock". It is locally so well cemented that it may rank as a calcrete. Late Pleistocene loess with a weakly developed soil commonly mantles gently sloping surfaces of irrigated farmland. Named by Malde and Powers (1962). Sadler and Link (1996) describe lithofacies and interpret provenance, paleocurrents, and age of the Tuana Gravel, which largely corroborate the descriptions of Malde and Powers (1962). The Tuana Gravel overlies Glenns Ferry Formation at the Hagerman Fossil Beds National Monument where the upper Glenns Ferry Formation records the base of the Kaena reverse polarity subchron with an estimated age of 3.11 Ma (Neville and others, 1979; Hart and Brueseke, 1999). The age of Tuana Gravel remains poorly constrained, but Malde (1991) and Othberg (1994) suggest the Tuana Gravel and the Tenmile Gravel near Boise, apparently graded to the same base level, are correlative. A minimum age for Tuana Gravel mapped to the northwest is 1.92 ± 0.16 Ma (Malde, 1991). A minimum age for the Tenmile Gravel is 1.58 ± 0.085 (Othberg, 1994). These gravels represent fluvial and glacial regimes driven by cooler climate in late Pliocene but before Pleistocene incision of the western Snake River Plain (Othberg, 1994).

Tsgf

**Glenns Ferry Formation (Pliocene)**—Poorly consolidated, bedded lake and stream deposits. In Hagerman Valley primarily flood plain lithofacies that include calcareous olive silt, dark clay, sand locally cemented, and fine-pebble gravel (Malde and Powers, 1962; Malde and Powers, 1972; Malde, 1972; McDonald and others, 1996). The formation includes intercalated but laterally inextensive beds of tephra and basalt flows (*Tsh*, for example). Repenning and others (1995) interpret the ages of various localities included in the Glenns Ferry Formation, and present a paleogeographic history of Pliocene to early Pleistocene lake and stream deposits in the western Snake River Plain. The basin-filling contribution of the Glenns Ferry Formation to the western Snake River Plain's tectonic subsidence is described by Wood and Clemens (2002). Mammalian fossils in deposits at the Hagerman Fossil Beds National Monument are Blancan (Pliocene) in age (Repenning and others, 1995). Hart and Brueseke (1999) corroborate the Pliocene age with Ar-Ar dates on intercalated basalt flows and ash beds ranging from 3.4 to 3.8 Ma.

## EOLIAN DEPOSITS

Qed

**Dune sand (Holocene)**—Stratified fine sand of stabilized wind dunes. Fills swales between basalt pressure ridges but sand is generally less than 4 feet thick. Agricultural modifications to the land have tended to smooth dune morphology.

## MASS MOVEMENT DEPOSITS

Qt

**Talus of Snake River Canyon walls (Holocene)**—Angular pebble-, cobble-, and boulder-size fragments of basalt that have broken off nearly vertical rock walls and accumulated below. Deposits are characterized by a steeply sloping surface that is at or near the angle of repose. Not mapped where thin talus partially covers older units.

Qlsa

**Deposits of active landslides (Holocene)**—Primarily unsorted and unstratified silt and clay in slumps, slides, and debris flows that originate in Glenns Ferry Formation, Yahoo Clay, or in older landslide deposits. On west side of Snake River new landslides have activated or old landslides have reactivated since 1979. The recent activity correlates with irrigated farming on the upland west of bluffs of the Snake River valley (Hagerman Fossil Beds National Monument, 2004). Aerial photographs flown in 1998 and 2004 were used to identify and map active landslides.

Qls **Landslide deposits (Holocene and Pleistocene)**—Unsorted and unstratified deposits comprised of fragments of sediments to broken clasts of basalt depending on location. The walls of Malad Gorge are formed of a complex of slumps composed of large basalt blocks and jumbled, angular cobbles and boulders of basalt. West of the Snake River, slumps, slides, and debris flows originating in the Glenns Ferry Formation are composed of unsorted and unstratified silt and clay. In the Snake River canyon north of Hagerman slumps and slides have displaced Yahoo Clay and basalt of McKinney Butte. The oldest landslides may have initiated during or just after the Bonneville Flood; others are probably Holocene. In addition to the landslide deposit, the unit includes the landslide scarp and the headwall (steep area adjacent to and below the landslide scarp) from which material broke away (see Symbols).

## BASALT UNITS

In the northeast part of the quadrangle Pleistocene basalt flows form the capping units along the east bluff of the Snake River canyon. They also occur as intracanyon flows on the valley floor west and north of Hagerman. Isolated erosional remnants of these basalts occur west of the Snake River. These lava flows were extruded from several shield volcanoes to the east and north. Nearly all of the basalt is slightly to very vesicular and most of the units are also diktytaxitic to some degree (i.e., containing voids with protruding crystals). Even units with a fine-grained groundmass commonly have a coarse, grainy texture. Older basalt flows are exposed at several locations along the Snake River and on the slopes west of the river intercalated with sediments of the Glenns Ferry Formation. These older units are generally grainy textured, less vesicular, and commonly altered. These characteristics cause the basalt to weather into fine granular detritus.

Qmk **Basalt of McKinney Butte (Pleistocene)**—Fine-grained basalt with very abundant plagioclase phenocrysts as crisscrossed interlocking crystals up to 1 cm in length, or as glomerocrysts commonly with olivine grains and clots. Voids are common among the interlocking crystals. The abundance of phenocrysts and the voids gives the rock a coarse-textured appearance. Remanent magnetic polarity is normal (Malde, 1971). Source is McKinney Butte located 15 miles north of Hagerman. Named McKinney Basalt by Stearns and others (1938); name retained by Malde (1971), Malde and Powers (1972) and Covington and Weaver (1990). Tauxe and others (2004) report an 40Ar/39Ar weighted mean plateau age of 0.052 Ma (their sample sr16, McKinney Basalt). Forms a thin capping unit north of Malad Gorge. Eolian sand (Qed) partly buries and smooths the unit's surface in the Hagerman quadrangle, but the sand diminishes 3 miles to the north where the basalt's surface is characterized by youthful flow features such as lava channels, tubes, collapse pits, and pressure ridges. Surface drainage is undeveloped.

Qnb **Basalt of Notch Butte (Pleistocene)**—Fine-grained, dark gray to black basalt with common to abundant olivine phenocrysts and clots 0.5 to 1.5 mm in diameter. Locally contains a few small plagioclase phenocrysts up to about 1 mm in length, or scarce to rare glomerocrysts of plagioclase and olivine up to about 5 mm in diameter. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is Notch Butte located 25 miles east of Hagerman. Several fingers of lava cascaded into the ancestral Snake River valley. Bonneville-Flood scoured remnants of the basalt are found west of Hagerman and at Lower Salmon Falls dam north of Hagerman. These areas were mapped as Sand Springs Basalt by Stearns and others (1938), Malde and others (1963), Malde and Powers (1972), and Covington and Weaver (1990), who identified the source as a shield volcano northeast of Twin Falls. However, based on paleomagnetic directions and chemical analyses, this Hagerman-Valley basalt probably originated from Notch Butte.

Qgb **Basalt of Gooding Butte (Pleistocene)**—Fine-grained basalt with scattered to abundant plagioclase phenocrysts up to 1 cm in length, and plagioclase-olivine intergrowths up to 1 cm in diameter. Olivine is olive-greenish-brown and is mostly clustered. The basalt is diktytaxitic and vesicular, with vesicles ranging from small and spherical to large and irregular. Common carbonate filling and coating in voids. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is Gooding Butte located 10 miles northeast of Hagerman. Outcrops uncommon on upland surface but well exposed in Malad Gorge and on the east Snake River canyon wall north of Hagerman. Equivalent to Thousand Springs Basalt, Malad Member of Malde and others (1963). Almost no basalt pressure ridges rise above a nearly complete mantle of loess, except where scoured by Malad River floods. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet; commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche is highly variable. Most of the land is cultivatable.

Qma **Basalt of Madson Spring (Pleistocene)**—Fine- to medium-grained, dark gray basalt with very abundant olivine grains and clots up to 4 mm in diameter. Exposed at one location west of the Snake River near the north edge of the quadrangle where it occurs as an erosional remnant resting on Glenns Ferry sediments. Equivalent to the Madson Basalt of Malde and Powers (1972). Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Tauxe and others (2004) report an $^{40}Ar/^{39}Ar$ weighted mean plateau age of 0.404 Ma for this unit (their sample sr12, Madson Basalt). Source undetermined, but Malde (1971) suggests the source was likely to the east. It may represent early eruptions from Gooding Butte.

Tsh

**Basalt of Shoestring Road (Pliocene)**—Basalt of Shoestring Road (Pliocene)—Medium-grained, gray to sooty brown basalt with common to abundant plagioclase laths as long as 1 cm and weathered olivine crystals. Remanent magnetic polarity is reverse, as determined in the field and as reported by Neville and others (1979). Forms a thin layer 30-50 feet thick within the Glenns Ferry sediments on the west side of the Snake River that extends northward from the Hagerman Horse Quarry (McDonald and others, 1996; Hart and Brueseke, 1999). Locally, the basalt flow thins, pinches out, or is replaced by layered volcaniclastic deposits. A thicker sequence of basalt is exposed in the northwest part of the quadrangle. Source may be vent of Tshv unit, or may be from an unidentified source to the north. Equivalent to the Shoestring Basalt of Malde and Powers (1972). May include the Deer Gulch basalt of Hart and Brueseke (1999) who dated the Shoestring basalt at 3.68 Ma and the Deer Gulch basalt at 3.40 Ma, although they report that both units occupy the same stratigraphic horizon in the Glenns Ferry Formation, and chemical compositions of the two units are very similar (Hart and Brueseke, 1999).

Tshv

**Basalt and vent material, basalt of Shoestring Road? (Pliocene)**—Basalt and vent-derived scoriaceous and ropy rubble, spatter, and basalt fragments. Basalt is fine grained and glassy in places with abundant plagioclase laths and glomerocrysts up to 1 cm long, and common to abundant olivine grains and clots 0.5-3 mm in diameter. Some olivine and plagioclase intergrowths. Invasive into Glenns Ferry Formation sediments, which are baked for several inches at the basalt-sediment contact. Malde and Powers (1972) considered this a vent for the Shoestring Basalt, but we did not confirm that correlation in the field and therefore have mapped it as a separate unit. Its position within the Glenns Ferry sediments, however, gives credence to their interpretation.

Tos

**Basalt of Oster Lakes (early Pliocene)**—Grainy, fine- to medium-grained but coarse-textured vesicular basalt. Dark gray to brownish gray or brick colored with a light purplish hue in places. Abundant plagioclase phenocrysts as much as 5 mm in length; groundmass is glassy in places. Unit is subaerial and quite fresh where exposed near the fish hatcheries at Oster Lakes. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Field relations suggest it stratigraphically and probably unconformably overlies the Tvd vent tephras and the altered basalt flows (*Tub*), which have reverse polarity where analyzed. Source or sources undetermined, but may be to the west or southwest. Previously mapped as "Banbury Basalt, basalt of upper part" by Malde and Powers (1972) and Covington and Weaver (1990). A K-Ar age determination by Armstrong and others (1975) on this unit is 4.4±0.6 Ma. An $^{40}Ar/^{39}Ar$ weighted mean age of 4.46±0.39 Ma was obtained on our sample 03VG039 (Esser, 2005).

Teh

**Basalt of Elmas Hill(Pliocene to late Miocene)**—Medium-grained abundantly plagioclase-phyric to dense, fine-grained aphyric basalt. Plagioclase-phyric basalt is very vesicular with a significant amount of caliche coating or filling in the vesicles. Some outcrops appear fresh while others are altered or degraded. Remanent magnetic polarity is reverse as determined in the field and from laboratory analysis. Named for butte just southwest of the map area with a survey point "Elmas" noted on the Crows Nest NE 7.5' topographic map. In the lower part of Yahoo Creek, Teh forms a gentle dip slope to about Crows Nest Road, where the unit steepens to about 15 degrees and dips beneath Glenns Ferry Formation sediments (*Tsgf*). Two flows are present in the gully where the dip steepens. The upper flow is dense, fine-grained and aphyric, but the older flow is degraded with sooty weathering and a more grainy texture. The chemical compositions of the two flows are dissimilar, indicating either chemical changes in the erupting magma or possibly different sources. Textural differences between plagioclase-phyric and aphyric exposures may be due to distance from the source or to separate eruptive events from one or more sources.

Tub

**Older basalt flows, undivided (early Pliocene to late Miocene**)—Medium- to coarse-grained, gray to sooty brown, dense to altered or weathered basalt flows exposed in the vicinity of Dolman Rapids and Dolman Island south of Hagerman, and at several locations along the Snake River north of Hagerman. Includes fine-grained sediments either intercalated with or underlying the basalt near the mouth of Malad River. Source(s) unknown but may be from the south and southwest. Age is poorly constrained, but underlies sediment of the Glenns Ferry Formation (*Tsgf*). All flows in this unit that were analyzed for remanent magnetism have reverse polarity, although all may not be age-equivalent and not all flows were analyzed. Malde and Powers (1972) included most of these flows in their "Banbury Basalt, basalt of upper part". Stearns and others (1938) included them in the Banbury volcanics.

## Appendix B

Survey Control

Coordinates and elevations were obtained using a Trimble GeoXT 2005 model GPS and occupying a point for a period to collect 100 readings which were post-processed in the office. The maximum PDOP value of 4 was selected to obtain the best accuracy this model is capable of. Quality control and assurance included using the same method and collection on a known US Geological Survey bench mark located in the Hagerman Valley at T7S R13E, section 15, SE of the SW. The bench mark has known coordinates from professional grade surveys and maintained by the National Geodetic Survey (http://www.ngs.noaa.gov/) and the data were compared against the GPS coordinates to obtain an absolute error value for evaluating the accuracy.

On the day of surveying the Rangen Monitor Well the absolute value for accuracy of the GPS, when evaluated against the bench mark was:

| | |
|---|---|
| easting | = 0.5 feet |
| northing | = 8.4 feet |
| elevation | = 0.9 feet |

On the day of surveying the Rangen Spring discharge the absolute value for accuracy was:

| | |
|---|---|
| easting | = 4.3 feet |
| northing | = 5.7 feet |
| elevation | = 3.6 feet |

The individual days of GPS data are consistent with the total running absolute value for the average accuracy for this same GPS unit during the past year which is:

| | |
|---|---|
| easting | = 3.4 feet |
| northing | = 5.4 feet |
| elevation | = 3.5 feet |

## References


Bonnichsen, B., and Breckenridge, R.M., 1982, Cenozoic Geology of Idaho, Idaho Geological Survey, Moscow, Idaho, Bulletin 26, 725 p.

Bjork, P.R., 1968, The carnivora of the Hagerman local fauna (late Pliocene) of southwestern Idaho, University of Michigan, Ph.D. dissertation, 165 p.

Cope, E.D., 1883, On the fishes of the Recent and Pliocene lakes of the western part of the Great Basin, and the Idaho Pliocene lake: Philadelphia Academy of Natural science Proceedings, v. 35, 134-166 p.

Covington, H.R. and J.N. Weaver, 1990, Geologic Map and Profiles of the North Wall of the Snake River Canyon, Bliss, Hagerman and Tuttle Quadrangles, Idaho: U.S. Geological Survey Miscellaneous Investigations Series Map MF-809.

Dallas, K., 2005, Hydrologic study of the Deer Gulch basalt in Hagerman fossil beds national monument, Idaho, thesis, 96p.

Erwin, F., 2008 personal communication.

Farmer, C.N., 2008, "Traffic jams occur in nature too" – geologic architecture of the Hagerman valley spring discharge areas, Idaho Water Resources Research Institute Ground Water Connections Conference, oral presentation.

Farmer, C.N., and Nagai, O., 2004, Non-native vegetation growth patterns as a tracer for development of human caused perched aquifers in landslide areas, Journal of the Japan Landslide Society, vol. 41, no. 3 (161), 12 p.

Farmer, C.N., and Larsen, I., 2001, Ground Water Tracer Tests at the Hagerman Fossil Beds National Monument, National Park Service, Technical Report.

Farmer, C.N., 1999, A hydrostratigraphic model for the perched aquifer systems located near the Hagerman fossil beds national monument; Idaho, thesis, 106 p.

Gillerman, V.S., J.D. Kauffman and K.L. Othberg, 2005, Geologic Map of the Thousand Springs Quadrangle, Gooding and Twin Falls Counties, Idaho: Idaho Geological Survey digital web map 49.

Kauffman, J.D., K.L. Othberg, V.S. Gillerman and D.L. Garwood, 2005a, Geologic Map of the Twin Falls 30 x 60 Minute Quadrangle, Idaho: Idaho Geological Survey digital web map 43.

Kauffman, J.D., K.L. Othberg and V.S. Gillerman, 2005b, Geologic Map of the Tuttle Quadrangle, Gooding and Twin Falls Counties, Idaho: Idaho Geological Survey digital web map 51.

Lee, D.E., 1995, Characterization of the Glenns Ferry Formation in the Fossil Gulch area, Hagerman Fossil Beds National Monument, Idaho: Geology Reports, Idaho Museum of Natural History, Geology Report #1, 59 p.

Lorkowski, R. M. and Hauser, J., 1996, Carbonaceous shale facies of the Glenn's Ferry Formation, Hagerman Fossil Beds National Monument, Idaho: National Park Service unpublished report.

Malde, H.E., 1991, Quaternary geology and structural history of the Snake River Plain, Idaho and Oregon in Morrison, R.B., ed., Quaternary nonglacial geology; conterminous U.S.: Boulder, CO, Geological Society of America, The Geology of North America, v. K-2.

Malde, H.E., 1972a, Geologic Map of the Glenns Ferry-Hagerman Area, west-central Snake River Plain, Idaho, U.S. Geological Survey Miscellaneous Investigations Map I-169, scale 1:48,000, 2 sheets.

Malde, H.E., 1972b, Stratigraphy of the Glenns Ferry Formation from Hammett to Hagerman, Idaho: U.S. Geological Survey Bulletin 1331-D, 19 p.

Malde, H.E., and Powers, H.A., 1962, Upper Cenozoic stratigraphy of western Snake River Plain, Idaho: Geological Society of America Bulletin, v. 73. no. 10, 1197-1220 p.

Malde, H.E., 1965, Snake river plain, *in* The Quaternary of the United States, A review volume for the VII congress of the international association for Quaternary research, Wright, H.E. and Frey D.G., editors, Princeton University Press, Princeton, New Jersey.

Malde, H.E., and Powers, H. A., 1958, Flood-plain origin of the Hagerman Lake Beds, Snake River Plain, Idaho (abs.): Geological Society of America Bulletin, v. 69, 1608 p.

Maley, T., 1987, Exploring Idaho geology: Mineral Land Publications, 232 p.

Othberg, K.L., V.S. Gillerman and J.D. Kauffman, 2005, Geologic Map of the Hagerman Quadrangle, Gooding and Twin Falls Counties, Idaho: Idaho Geological Survey digital web map 50.

Saddler, J.L., 1997, Sedimentology and stratigraphy of the Pleistocene Tuana gravel at Hagerman Fossil Beds National Monument, Idaho: Idaho State University, M.S. thesis, 62 p.

United States Bureau of Reclamation, 2001, Water measurement manual, http://www.usbr.gov/pmts/hydraulics_lab/pubs/wmm/.

Welhan, J., 2008, Inland northwest research alliance subsurface science graduate program; subsurface architecture field trip, handbook 43 p.